



Contents lists available at ScienceDirect

Developmental and Comparative Immunology

journal homepage: www.elsevier.com/locate/dci

Short communication

Limiting immunopathology: Interaction between carotenoids and enzymatic antioxidant defences

A. Babin^{*}, C. Siciat, M. Teixeira, J.-P. Troussard, S. Motreuil, J. Moreau, Y. Moret

Équipe Écologie Évolutive, UMR CNRS 6282 Biogéosciences, Université de Bourgogne, Dijon, France

ARTICLE INFO

Article history:

Received 29 August 2014

Revised 9 December 2014

Accepted 10 December 2014

Available online 15 December 2014

Keywords:

Carotenoids

Antioxidant

Immunopathology

Crustacean

Ecological immunology

Immuno-stimulants

ABSTRACT

The release of reactive oxygen and nitrogen species (ROS and RNS) during the inflammatory response generates damages to host tissues, referred to as immunopathology, and is an important factor in ecological immunology. The integrated antioxidant system, comprising endogenous antioxidant enzymes (e.g. superoxide dismutase SOD, and catalase CAT) and dietary antioxidants (e.g. carotenoids), helps to cope with immune-mediated oxidative stress. Crustaceans store large amounts of dietary carotenoids for yet unclear reasons. While being immunostimulants and antioxidants, the interaction of these pigments with antioxidant enzymes remains unclear. Here, we tested the interaction between dietary supplementation with carotenoids and immune challenge on immune defences and the activity of the antioxidant enzymes SOD and CAT, in the amphipod crustacean *Gammarus pulex*. Dietary supplementation increased the concentrations of circulating carotenoids and haemocytes in the haemolymph, while the immune response induced the consumption of circulating carotenoids and a drop of haemocyte density. Interestingly, supplemented gammarids exhibited down-regulated SOD activity but high CAT activity compared to control ones. Our study reveals specific interactions of dietary carotenoids with endogenous antioxidant enzymes, and further underlines the potential importance of carotenoids in the evolution of immunity and/or of antioxidant mechanisms in crustaceans.

© 2014 Elsevier Ltd. All rights reserved.

1. Introduction

During the inflammatory response, the release of highly reactive oxygen and nitrogen species (ROS and RNS) generates damages to host tissues, referred to as immunopathology, and contributes to immune system costs uncovered in evolutionary studies (Pursall and Rolff, 2011). Among the protective processes limiting inflammatory costs, hosts evolved the antioxidant defence system controlling the amount of circulating cytotoxic ROS and RNS. This system comprises key endogenous antioxidant enzymes, including the superoxide dismutase (SOD) and catalase (CAT), which are successively involved in the detoxification of superoxide radicals, and the action of dietary antioxidants such as carotenoids (Felton and Summers, 1995). While immunostimulants (e.g. Park et al., 2010), carotenoids also scavenge free radicals (e.g. ROS and RNS), which may limit the costs of inflammation (e.g. Walrand et al., 2005).

Despite their involvement in the antioxidant defence system, the interaction of dietary carotenoids with the key endogenous antioxidant enzymes SOD and CAT remains unclear. On the one hand, carotenoids may take over the action of the antioxidant enzymes

(Chien et al., 2003; Wang et al., 2006), reducing their relative activity and therefore their maintenance costs. On the other hand, in addition of being free radical scavengers, carotenoids were suggested to stimulate the activity of these antioxidant enzymes (Kim et al., 2005; Lee et al., 2011), which would enhance detoxification efficacy upon immune activity. In both cases, carotenoids may improve the detoxification of free radicals, alleviating the costs associated with oxidative stress. This might be particularly relevant for animals storing large amounts of carotenoids in their tissues.

Crustaceans store naturally large amounts of dietary carotenoids, mainly astaxanthin and lutein, in their tissues (Dembitsky and Rezanka, 1996; Gaillard et al., 2004) for yet unclear reasons. In the amphipod, *Gammarus pulex*, the amount of carotenoids in the haemolymph varies among natural populations and correlates positively with immune defences (Cornet et al., 2007). Crustacean immunity is innate and relies on the response of immune cells and rapidly activated enzyme cascades such as the prophenoloxidase (proPO) that are at the core of the inflammatory response (Nappi and Ottaviani, 2000; Söderhäll and Cerenius, 1998). Interestingly, the experimental supplementation of the food of *G. pulex* with carotenoids broadly stimulated innate immune defences without increasing the inflammation cost (Babin et al., 2010). Beneficial effects of dietary carotenoids combined to the high capacity of crustaceans to store these pigments suggest that the evolution of crustacean immunity may occur in a particular context. By

^{*} Corresponding author. UMR CNRS 6282 Biogéosciences, Université de Bourgogne, 6 Boulevard Gabriel, Dijon, France. Tel.: +33380399024; fax.: +33380396231.

E-mail address: aurelie.babin@hotmail.fr (A. Babin).

reducing inflammation-associated costs, carotenoids may alleviate part of this constraint on the evolution of the immune function (Babin et al., 2010). To better understand the potential importance of carotenoids in the evolution of immunity and/or of antioxidant mechanisms in crustaceans, we aimed to test whether the action of these pigments stored by crustaceans is additive or synergistic with that of endogenous antioxidant enzymes to reduce immunopathology.

To this purpose, we tested the influence of food supplementation with carotenoids on important cellular and humoral components of the immune system of *G. pulex* (density of haemocytes, activity of the proPO system) and on the activity of the key endogenous antioxidant enzymes SOD and CAT, after a bacterial immune challenge.

2. Methods

2.1. Experimental design and measures

One thousand male gammarids were collected in March 2011 in the river Suzon (Val-Suzon, eastern France), and maintained individually in glass vials with 20 mL of oxygenated dechlorinated UV-treated tap water under standard laboratory conditions ($15\text{ }^{\circ}\text{C} \pm 1\text{ }^{\circ}\text{C}$, 12:12 light:dark cycle). Immediately after field collection, gammarids were starved for 3 days before exposure to the food treatment to exacerbate hunger (Babin et al., 2010). For the food treatment, 500 gammarids were supplied with food supplemented with carotenoids (carotenoid food treatment) whereas the other 500 were offered the same food basis as in the carotenoid food treatment but without carotenoids (control food treatment). The food recipe is identical to the one used by Babin et al. (2010). The food treatment lasted 17 days during which gammarids were provided daily with food ad libitum and the remains from the previous provision were removed to keep the containers clean. Survival was checked daily and water was changed once a week to provide sufficient oxygen to the gammarids.

After the food treatment, 150 supplemented and 150 non-supplemented gammarids were randomly assigned to one of the three immune treatments: 'naive', 'PBS' and 'bacteria' to test interactions between immune challenge and carotenoid supplementation on carotenoid concentration, haemocyte density, activities of the proPO system in their haemolymph, and activity of the SOD and CAT enzymes of their antioxidant system from their entire body.

The 'naive' group was unchallenged. The 'PBS' group received a single injection of 0.5 μL of sterile phosphate buffer saline (PBS: 8.74 g NaCl; 1.78 g Na_2HPO_4 , $2\text{H}_2\text{O}$; 1000 mL of distilled water; pH 6.5). The 'bacteria' group received an 8×10^5 cells. mL^{-1} dose of heat-killed *Escherichia coli* (CIP 103410, Institut Pasteur, Paris, France; inactivated 5 min at $100\text{ }^{\circ}\text{C}$) in 0.5 μL of PBS. While non-pathogenic, heat killed-bacteria stimulate the innate immune response, leading to the production of cytotoxic ROS and RNS in invertebrates (Nappi and Ottaviani, 2000), and promote immunopathology (Sadd and Siva-Jothy, 2006). Bacterial growth and injection procedures were done as described in Cornet et al. (2009). After injection, gammarids were maintained individually under standard conditions without food for 15 hours before haemolymph sampling. This is the time required for the immune response of *G. pulex* to complete bacterial clearance and restore homeostasis (Moret, personal communication).

Haemolymph was collected as described in Cornet et al. (2007). Each gammarid provided up to 3 μL of haemolymph, immediately diluted in 20 μL of ice-cold PBS, vortexed and split into a 10 μL sample for carotenoid dosage and a 13 μL sample for immune parameters, either haemocyte density or activities of the proPO system. Samples allocated to measure carotenoid concentration and activities of the proPO enzyme, as well as individuals, were frozen in liquid nitrogen and stored at $-80\text{ }^{\circ}\text{C}$ until later examination. Samples al-

located to measure density of haemocytes were immediately treated using a Neubauer-improved haemocytometer under a microscope (magnification $\times 400$).

Carotenoids were extracted and quantified via a colorimetric assay following the method of Cornet et al. (2007). Values were corrected to obtain concentrations for 1 μL of pure haemolymph. The activities of naturally activated PO (PO activity) and that of the pro-enzymes (which give the total PO activity when added together) were measured using a spectrophotometric assay following Cornet et al. (2009). Enzyme activity was measured as the slope (V_{max} value in milli-absorbance. min^{-1}) of the reaction curve using the software SOFT-Max@Pro 4.0 (Molecular Devices), and reported as the activity of 1 μL of pure haemolymph.

SOD and CAT enzyme activities were scored spectrophotometrically following the protocol of De Block and Stoks (2008). SOD activity was quantified based on the protocol from the SOD assay kit from Cayman Chemical (Ann Arbor, MI, USA), which measures the formation of formazan dye upon reduction of tetrazolium salt with superoxide anions. SOD activity was expressed in units (U) of SOD per milligram of gammarid fresh mass (U/mg FM). One absorbance unit of SOD corresponds to the amount of enzyme needed to inhibit the colorimetric reaction by 50%. CAT activity was quantified following a protocol adapted from Aebi (1984). We measured the removal of the H_2O_2 as the reduction in absorbance at 240 nm within 3 min at $25\text{ }^{\circ}\text{C}$. CAT activity was expressed as the decline in absorbance units (U) per min and per mg of gammarid fresh mass (U/min/mg FM) (see details in Supplementary material). Because we aimed to examine variation at the level of the whole organism and compare individuals, the measurements discussed earlier are expressed per unit of volume of haemolymph (e.g. concentration of carotenoids, density of haemocytes and activity of the proPO system) or unit of fresh mass (e.g. activity of the SOD and the CAT enzymes).

2.2. Statistics

Gammarid survival during the first 17 days of food treatment was analysed using a chi-square test. The effect of the immune challenge on carotenoid concentration, PO activity, total PO activity, density of haemocytes, SOD activity and CAT activity in interaction with the food treatment, was analysed using a linear model for each dependent variable including gammarid fresh mass as covariate, food treatment and immune treatment as fixed factors, and their two-way interactions. Non-significant interaction terms were removed from the models. Data of the six dependent variables were square root transformed to homogenise the variance. All data were analysed using SPSS 19 for Macintosh.

3. Results/discussion

Over the 17 days of food treatment, dietary supplementation with carotenoids significantly improved survival (75.2% versus 60.8%, $\chi^2_{(1)} = 23.17$, $P < 0.001$). Supplemented gammarids had higher carotenoid concentrations in their haemolymph (Fig. 1A; Table 1) within the natural range (up to 1840 ng/ μL ; Cornet et al., 2007) and exhibited stronger reddish coloration (Supplementary Fig. S1). The immune treatment affected carotenoid concentration in interaction with the food treatment (Table 1). Whereas the carotenoid concentration of supplemented gammarids remained high across immune treatments, the immune challenge with heat-killed bacteria consumed a significant amount of these circulating pigments in non-supplemented ones (Fig. 1A). This clearly illustrates the physiological involvement of carotenoids in the immune response, probably by scavenging the excess of free radicals produced by immune activity (Walrand et al., 2005). Saturation of the haemolymph with carotenoids in supplemented gammarids may




Fig. 1. Effect of the 17 days of food supplementation with carotenoids (control, open circles; supplemented, filled triangles) and the immune treatment (naive, PBS, heat-killed bacteria) on (A) the concentration of circulating carotenoids (in ng/μL), (B) the density of haemocytes (in cells/μL), and (C, D) the activity of the endogenous antioxidant enzymes SOD (C, in U/mg fresh mass) and CAT (D, U/min/mg fresh mass) of *Gammarus pulex*. Error bars are standard errors of means.

have compensated for the immune-associated depletion of carotenoid contents. The influence of the food treatment on carotenoid concentration depended also on gammarid's body mass (Table 1). Carotenoid concentration tended to co-vary negatively with body mass of supplemented gammarids ($R^2 = 0.03$, $F_{1,136} = 3.72$ $P = 0.056$), whereas it co-varied positively with body mass of non-supplemented gammarids ($R^2 = 0.09$, $F_{1,132} = 13.65$ $P < 0.001$).

Dietary supplementation enhanced the basal concentration of haemocytes (Fig. 1B; Table 1), confirming the immunostimulant effect of these pigments (Babin et al., 2010). Regardless of the food treatment, haemocyte density dropped after the immune challenge with heat-killed bacteria (Fig. 1B; Table 1), demonstrating the occurrence of an immune response. Upon infection, circulating haemocytes disappear as they are recruited for phagocytosis (Jiravanichpaisal et al., 2006; Söderhäll et al., 2003), which goes along with the oxidative burst resulting from the release of superoxide radicals (Nappi and Ottaviani, 2000). Variation in PO and total PO activity was only explained by gammarids' body mass, with lower levels in heavy animals (Table 1; $R^2 = 0.12$, $F_{1,137} = 19.12$ $P < 0.001$; total PO: $R^2 = 0.13$, $F_{1,137} = 20.10$ $P < 0.001$). Despite involved in defence against bacterial infections (Söderhäll and Cerenius, 1998), the activity of the proPO system does not always predict bacterial clearance in *G. pulex* (Cornet et al., 2009) and in insects (Adamo, 2004; Ayres and Schneider, 2008; Binggeli et al., 2014).

SOD and CAT activities are typically expected to increase when stress conditions increase metabolic rate. Here, the activity of both enzymes depended on the interaction between the immune response and dietary supplementation (Table 1). These two enzymes act successively on superoxide radicals: the SOD converts super-

oxide into hydrogen peroxide, which is then detoxified by the CAT into water and oxygen (Felton and Summers, 1995). Supplemented gammarids exhibited relatively low but similar levels of SOD activity across immune treatments compared to that of non-supplemented gammarids. The latter ones exhibited relatively high enzymatic activity, except for the drop of SOD activity at a similar level to that of supplemented gammarids after the immune challenge with heat-killed bacteria (Fig. 1C). This suggests that stored astaxanthin in supplemented gammarids have acted as free radical scavengers (Fassett and Coombes, 2011), taking over SOD enzymes and lowering their activity for detoxification of superoxide radicals likely produced during basal metabolic activity. Unexpectedly, the immune challenge caused a drop of SOD activity in non-supplemented gammarids (Fig. 1C). As the immune response to a systemic bacterial challenge is energetically costly (Moret and Schmid-Hempel, 2000) in addition to releasing superoxide radicals into the haemocoel (Nappi and Ottaviani, 2000), high level of SOD activity could probably not be sustained simultaneously with the immune response. However, this hypothesis appears unlikely because a similar reduction of CAT enzyme activity would be expected (but see Fig. 1D). Nevertheless, the drop of SOD activity in non-supplemented gammarids mirrors that of circulating carotenoids in the haemolymph (Fig. 1A, C). Free radicals released by the immune response might have been scavenged by the small amount of circulating carotenoids, reducing their concentration and SOD activity. Besides, the influence of the carotenoids supplementation on SOD activity also depended on gammarids' body mass (Table 1), with lower SOD activity in heavier animals for non-supplemented gammarids only.

Table 1

Univariate models testing the effect of food supplementation with carotenoids ('Food'), immune treatment ('Treatment') and body mass ('Mass') on carotenoid concentration, levels of immune defences (haemocyte density, PO and total PO activities) and the activity of endogenous antioxidant enzymes (SOD and CAT) of *Gammarus pulex*. The table provides the Fisher-test value (F), corresponding degrees of freedom (d.f.) and the probability (P) of obtaining the observed sample results under the null hypothesis condition (Zar, 1999). Note that half of the gammarids sampled per food treatment and challenge treatment were used to measure their density of haemocytes and the other half for the activity of the pro-PO system, explaining variation in degrees of freedom across dependent variables.

Dependent variables	Sources of variation	F	d.f.	P
Carotenoids	Model	16.96	7,264	<0.001
	Food	38.90	1,264	<0.001
	Treatment	1.99	2,264	0.138
	Mass	2.93	1,264	0.088
	Food × Treatment	4.61	2,264	0.011
Haemocytes	Food × Mass	17.13	1,264	<0.001
	Model	3.09	4,126	0.018
	Food	0.00	1,126	0.002
	Treatment	6.15	2,126	0.003
	Mass	0.47	1,126	0.496
PO	Model	4.83	4,134	0.001
	Food	0.00	1,134	0.973
	Treatment	0.27	2,134	0.762
	Mass	18.03	1,134	<0.001
	Total-PO	Model	6.09	4,134
SOD	Food	1.39	1,134	0.240
	Treatment	1.22	2,134	0.299
	Mass	19.26	1,134	<0.001
	Model	9.87	7,265	<0.001
	Food	10.38	1,265	0.001
CAT	Treatment	6.69	2,265	0.001
	Mass	2.06	1,265	0.152
	Food × Treatment	4.56	2,265	0.011
	Food × Mass	6.58	1,265	0.011
	Model	26.15	6,266	<0.001
CAT	Food	55.96	1,266	<0.001
	Treatment	0.17	2,266	0.850
	Mass	85.53	1,266	<0.001
	Food × Treatment	10.94	2,266	<0.001

As for SOD activity, CAT activity depended on the interaction between the immune and the food treatments (Table 1). However, the immune challenge with heat-killed bacteria did not reduce CAT activity (Fig 1D), but on the opposite promoted it, especially in supplemented gammarids. Due to its chemical structure consisting of hydroxyl groups at both ends of a carbon chain (Fassett and Coombes, 2011), astaxanthin may chemically interact with superoxide radicals and release hydrogen peroxides by yet unknown redox reaction, which are then detoxified by CAT activity. This may explain low SOD activity upon immune challenge in supplemented gammarids whereas CAT activity was higher and even stimulated by the immune response (Fig. 1C, D). The specificity of the interaction between dietary carotenoids and SOD activity might explain that the drop of carotenoid concentration in non-supplemented gammarids (Fig. 1A) did not impact CAT activity. As for PO and total PO activities, CAT activity was lower in heavier individuals (Table 1; $R^2 = 0.19$ $F_{1,271} = 63.56$ $P < 0.001$).

To summarise, dietary supplementation with carotenoids increased the concentrations of circulating carotenoids and haemocytes in *G. pulex*. The immune response consumed carotenoids, which are likely used to scavenge immune-released free radicals. Moreover, stored carotenoids likely took over the SOD activity, acting first in the detoxification chain of superoxide radicals, resulting in its reduced activity, whereas they promoted the CAT activity, acting in second in the detoxification chain. While these results demonstrate a specific interaction between dietary carotenoids and enzymes of the antioxidant defence system of *G. pulex*, they suggest that crus-

taceans may have a strong advantage of storing dietary carotenoids to limit immunopathology under high infection prevalence.

Acknowledgements

DSM, France, kindly provided Carophyll Pink 10%. AquaTechna, France, kindly provided samples of premix Vitatech-S. This work was supported by the CNRS and the ANR (ANR-08-JCJC-0006).

Appendix: Supplementary material

Supplementary data to this article can be found online at doi:10.1016/j.dci.2014.12.007.

References

- Adamo, S.A., 2004. How should behavioural ecologists interpret measurements of immunity? *Anim. Behav.* 68, 1443–1449.
- Aebi, H., 1984. Catalase in vitro. *Method Enzymol.* 105, 121–126.
- Ayres, J.S., Schneider, D.S., 2008. A signalling protease required for melanization in *Drosophila* affects resistance and tolerance of infections. *PLoS Biol.* 6 (12), e305.
- Babin, A., Biard, C., Moret, Y., 2010. Dietary supplementation with carotenoids improves immunity without increasing its cost in a crustacean. *Am. Nat.* 176, 234–241.
- Binggeli, O., Neyen, C., Poidevin, M., Lemaitre, B., 2014. Prophenoloxidase activation is required for survival to microbial infections in *Drosophila*. *PLoS Pathog.* 10 (5), e1004067.
- Chien, Y.H., Pan, C.H., Hunter, B., 2003. The resistance to physical stresses by *Penaeus monodon* juveniles fed diets supplemented with astaxanthin. *Aquaculture* 216, 177–191.
- Cornet, S., Biard, C., Moret, Y., 2007. Is there a role for antioxidant carotenoids in limiting self-harming immune response in invertebrates? *Biol. Lett.* 3, 284–288.
- Cornet, S., Biard, C., Moret, Y., 2009. Variation in immune defence among populations of *Gammarus pulex* (Crustacea: Amphipoda). *Oecologia* 159, 257–269.
- De Block, M., Stoks, R., 2008. Compensatory growth and oxidative stress in a damselfly. *Proc. R. Soc. B* 275, 781–785.
- Dembitsky, V.M., Rezanka, T., 1996. Comparative study of the endemic freshwater fauna of Lake Baikal. 7. Carotenoid composition of the deep-water amphipod crustacean *Acanthogammarus* (Brachyropus) *grewingkii*. *Comp. Biochem. Phys.* B 114, 383–387.
- Fassett, R.G., Coombes, J.S., 2011. Astaxanthin: a potential therapeutic agent in cardiovascular disease. *Marine Drugs* 9, 447–465.
- Felton, G.W., Summers, C.B., 1995. Antioxidant systems in insects. *Arch. Insect Biochem.* 29, 187–197.
- Gaillard, M., Juillet, C., Cezilly, F., Perrot-Minnot, M.J., 2004. Carotenoids of two freshwater amphipod species (*Gammarus pulex* and *G. roeseli*) and their common acanthocephalan parasite *Polymorphus minutus*. *Comp. Biochem. Phys.* B 139, 129–136.
- Jiravanichpaisal, P., Lee, B.L., Soderhall, K., 2006. Cell-mediated immunity in arthropods: hematopoiesis, coagulation, melanization and opsonization. *Immunobiology* 211, 213–236.
- Kim, J.H., Kim, Y.S., Song, G.G., Park, J.J., Chang, H.I., 2005. Protective effect of astaxanthin on naproxen-induced gastric antral ulceration in rats. *Eur. J. Pharmacol.* 514, 53–59.
- Lee, D.H., Kim, C.S., Lee, Y.J., 2011. Astaxanthin protects against MPTP/MPP plus -induced mitochondrial dysfunction and ROS production in vivo and in vitro. *Food Chem. Toxicol.* 49, 271–280.
- Moret, Y., Schmid-Hempel, P., 2000. Survival for immunity: the price of immune system activation for bumblebee workers. *Science* 290, 1166–1168.
- Nappi, A.J., Ottaviani, E., 2000. Cytotoxicity and cytotoxic molecules in invertebrates. *Bioessays* 22, 469–480.
- Park, J.S., Chyun, J.H., Kim, Y.K., Line, L.L., Chew, B.P., 2010. Astaxanthin decreased oxidative stress and inflammation and enhanced immune response in humans. *Nutr. Metab.* 7.
- Pursall, E.R., Rolff, J., 2011. Immune responses accelerate ageing: proof-of-principle in an insect model. *PLoS ONE* 6.
- Sadd, B.M., Siva-Jothy, M.T., 2006. Self-harm caused by an insect's innate immunity. *Proc. R. Soc. B* 273, 2571–2574.
- Söderhäll, I., Bangyeekhun, E., Mayo, S., Soderhall, K., 2003. Hemocyte production and maturation in an invertebrate animal; proliferation and gene expression in hematopoietic stem cells of *Pacifastacus leniusculus*. *Dev. Comp. Immunol.* 27, 661–672.
- Söderhäll, K., Cerenius, L., 1998. Role of the prophenoloxidase-activating system in invertebrate immunity. *Curr. Opin. Immunol.* 10, 23–28.
- Walrand, S., Farges, M.C., Dehaese, O., Cardinault, N., Minet-Quinard, R., Grolier, P., et al., 2005. In vivo and in vitro evidences that carotenoids could modulate the neutrophil respiratory burst during dietary manipulation. *Eur. J. Nutr.* 44, 114–120.
- Wang, Y.J., Chien, Y.H., Pan, C.H., 2006. Effects of dietary supplementation of carotenoids on survival, growth, pigmentation, and antioxidant capacity of characins, *Hyphessobrycon callistus*. *Aquaculture* 261, 641–648.
- Zar, J.H., 1999. *Biostatistical Analysis*, 4th ed. Prentice-Hall Inc.